

# Evaluation of Apramycin Activity Against Methicillin-Resistant, Methicillin-Sensitive, and Vancomycin-Intermediate *Staphylococcus aureus* Clinical Isolates

Katherine A. Truelson<sup>1</sup>, Thea Brennan-Krohn<sup>2</sup>, Kenneth P. Smith<sup>2</sup>, and James E. Kirby<sup>2,\*</sup>

<sup>1</sup>Boston University, Boston, MA, USA

<sup>2</sup>Department of Pathology, Beth Israel Deaconess Medical Center, Boston, MA, USA

Running title: Apramycin Activity vs. MRSA, MSSA, and VISA

Word count abstract:

Word count body of text:

#Corresponding Author

James E. Kirby

Beth Israel Deaconess Medical Center

330 Brookline Avenue - YA309

Boston, MA 02215

[jekirby@bidmc.harvard.edu](mailto:jekirby@bidmc.harvard.edu)

Phone: 617-667-3648

Fax: 617-667-4533

## Abstract

We evaluated the *in vitro* activity of apramycin against clinical strains of vancomycin-intermediate and methicillin-resistant and -susceptible *Staphylococcus aureus*. Apramycin demonstrated an MIC<sub>50</sub>/MIC<sub>90</sub> of 8/16 µg/mL. No strains had an MIC above the epidemiological cutoff value of 32 µg/mL, suggesting apramycin resistance mechanisms are rare in this strain population. The mounting evidence for broad-spectrum *in vitro* activity of apramycin against *S. aureus* and other bacterial species suggests that further exploration of apramycin or derivatives as repurposed human therapeutics is warranted.

## Keywords

apramycin, *Staphylococcus aureus*, activity spectrum, repurposing, antibiotic, natural product, synergy, aminoglycoside, gentamicin

## Abbreviations

## 1. Introduction

*Staphylococcus aureus* is both a human skin commensal and an opportunistic pathogen. It is the leading cause of bacteremia and infective endocarditis, as well skin and soft tissue, osteoarticular, and surgical site infections (Akhi, et al., 2017; Deleo, et al., 2010; Tong, et al., 2015). Strains resistant to methicillin and by proxy, all  $\beta$ -lactams (so called methicillin resistant *S. aureus* or MRSA), are common in the United States (Tong, et al., 2015). Vancomycin is the first-line treatment for methicillin-resistant *S. aureus* (MRSA) infection. However, limitations of vancomycin include relatively lower bactericidal activity compared with  $\beta$ -lactams; nephrotoxicity, associated with high dosages and underlying risk factors; and a requirement for routine monitoring of drug levels to ensure adequate dosing (Hazlewood, et al., 2010). Furthermore, vancomycin tolerant and heteroresistant populations may emerge during treatment (Bamberger & Boyd, 2005; Dombrowski & Winston, 2008; Hawkins, et al., 2007; Lodise, et al., 2008; Steinkraus, et al., 2007).

Therefore, additional treatment options for *S. aureus* would be welcome, especially for those strains with resistance to newer agents such as daptomycin and linezolid. aminoglycosides are not currently used to treat *S. aureus* infections as single agents. In particular, renal and ototoxic side effects are of significant concern (Jose, et al., 2010; Matt, et al., 2012). Therefore aminoglycoside therapy is generally reserved for treatment of Gram-negative infection, where pharmacodynamic considerations are considered more favorable (Tam, et al., 2006). Currently, gentamicin treatment of *S. aureus* is advocated only in low dose (1mg/kg q8 h or 3-5mg/kg q 24h) in combination with vancomycin or a  $\beta$ -lactam and rifampin, during therapy of staphylococcal endocarditis where prosthetic material is present and in combination with daptomycin for persistent bacteremia (Liu, et al., 2011). However, the relative benefit of gentamicin adjunctive therapy versus risk of kidney damage has been a subject of extensive debate (Bruss, 2009; Buchholtz, et al., 2009; Buchholtz, et al., 2011; Cosgrove, et al., 2009;

Frippiat, et al., 2009). Therefore, a non-toxic aminoglycoside with predictable activity against *S. aureus* would therefore be especially welcome.

Apramycin is a structurally unique aminoglycoside used in veterinary medicine and is characterized by a bicyclic sugar moiety and a 4-monosubstituted 2-deoxystreptamine ring (Meyer, et al., 2014). In contrast to aminoglycosides currently used in human therapy, apramycin appears highly selective for bacterial ribosomes and on this basis thought devoid of significant ototoxic and renal toxic side effects (Matt, et al., 2012). Previous data from our group and others have shown broad-spectrum *in vitro* activity of apramycin against carbapenem-resistant Enterobacteriaceae and multidrug-resistant *Acinetobacter baumannii* and *Pseudomonas aeruginosa* (Kang, et al., 2017; Livermore, et al., 2011; K.P. Smith & J.E. Kirby, 2016a, 2016b). Furthermore, we recently demonstrated potent *in vivo* bactericidal activity against *Acinetobacter baumannii* in a murine thigh infection model (Kang, et al., 2018).

However, activity spectrum data for *S. aureus* is limited. Therefore, the goal of this study was to explore *in vitro* activity of apramycin against contemporary clinical isolates of *S. aureus*.

## 2. Materials and Methods

### 2.1 Bacterial Strains and Antimicrobials

We evaluated a collection of 109 strains of *S. aureus* for their susceptibility to apramycin. Fourteen *S. aureus* strains obtained from the FDA-CDC Antimicrobial Resistance Isolate Bank (<https://www.cdc.gov/drugresistance/resistance-bank/>) were part of the vancomycin intermediate *S. aureus* panel with vancomycin MIC values of 4-8  $\mu\text{g}/\text{mL}$ . 95 additional strains of de-identified *S. aureus* clinical isolates were obtained from the Beth Israel Deaconess Medical Center (Boston, MA) clinical microbiology laboratory. Our strain collection was comprised of 38.5% methicillin-resistant *S. aureus* (MRSA), 48.6% methicillin-susceptible *S. aureus* (MSSA) and 12.9% vancomycin-intermediate *S. aureus* (VISA). All strains were stored frozen at  $-80^{\circ}\text{C}$  in a stock

solution of 50% glycerol and 50% cation-adjusted Mueller-Hinton broth (BD Diagnostics, Franklin Lakes, NJ) until use in experiments.

Apramycin sulfate was obtained from Alfa Aesar (Tewksbury, MA) and was dissolved in deionized water at 32 mg/ml and stored at -20°C in aliquots that were used only once. For synergy studies, apramycin, gentamicin, daptomycin, and vancomycin were dissolved at 100, 100, 10, and 10 mg/ml in water supplemented with 0.3% polysorbate 20 (P-20; Sigma–Aldrich), respectively. Linezolid was dissolved at 30 mg/ml in DMSO (Sigma-Aldrich). Solvents used reflect requirements for liquid handling by the HP D300 Digital Dispensing System (HP Inc, Palo Alto, CA) (Brennan-Krohn, et al., 2017; K. P. Smith & J. E. Kirby, 2016).

## 2.2 Susceptibility Testing

The Clinical Laboratory and Standards Institute (CLSI) broth microdilution reference method was used for MIC testing of apramycin (CLSI, 2015). MIC panels were created by serial dilution of stock apramycin with cation-adjusted Mueller-Hinton broth in round bottom, 96-well plates (Evergreen Scientific, Los Angeles, CA). Dilutions were prepared at 2x the final concentration in volumes of 50  $\mu$ L with final concentrations ranging from 1-256  $\mu$ g/mL after an equal volume of bacterial inoculum ( $5 \times 10^5$  cfu/ml final concentration) was added. Stock solutions were quality controlled against *S. aureus* ATCC 29213 on three separate days. All MIC values for ATCC 29213 were consistently 4  $\mu$ g/ml, which was in the middle of the 2-8  $\mu$ g/mL acceptable quality control range suggested by CLSI. Bacterial inocula were prepared by passaging previously frozen bacterial strains on trypticase soy agar containing 5% sheep's blood at 37°C. Isolated colonies were then suspended in cation-adjusted Mueller-Hinton broth for a final inoculum concentration of  $5 \times 10^5$  CFU/mL. After inoculation, broth microdilution plates were incubated at 35°C in ambient air for 16-20 hours. Each experiment also included both a positive (*S. aureus* ATCC 25923) and a negative control to which no organisms were added. *S. aureus* 25923 always showed an MIC of 4 or 8  $\mu$ g/ml supporting consistency of assay readout.

### ***2.3 MIC Interpretation***

The MIC of each strain from our collection was determined in duplicate on separate days. If duplicate MICs were within one doubling dilution of each other but were not the same, the higher MIC was used. If duplicate MICs were not within one doubling dilution of each other, a third replicate was performed, and the MIC was defined as the modal MIC of the three replicates. Categorical breakpoints for apramycin are not available either from the CLSI or the European Committee for Antimicrobial Susceptibility Testing and, therefore, categorical interpretation was not performed.

### ***2.4 Time-Kill Studies***

Time-kill studies were performed according to CLSI guidelines (Clinical and Laboratory Standards Institute, 1999). To prepare a starting inoculum for the time-kill studies, 100  $\mu$ L of a 0.5 McFarland suspension of colonies from an overnight plate was added to 5 mL of CAMHB and incubated on a shaker in ambient air at 35°C until it reached log phase growth. The log phase culture was then adjusted to a turbidity of 1.0 McFarland in CAMHB, and 100  $\mu$ L of this suspension was added to 10 mL volumes of antimicrobial solutions in cation-adjusted Mueller-Hinton broth (BD Diagnostics, Franklin Lakes, NJ) at time 0. Antibiotic concentrations selected were based on multiples of each isolate's MIC as determined through broth microdilution assays. A growth control and a negative control were run in parallel with each experiment. Cultures were incubated on a shaker in ambient air at 35°C.

Aliquots from the culture were removed at 0, 1, 2, 4, 6, and 24 hours. A serial 10-fold dilution in 0.9% sodium chloride was prepared and a 10  $\mu$ L drop from each dilution was transferred to a Mueller Hinton plate (Thermo Fisher, Waltham, MA) (Chen, et al., 2003; Herigstad, et al., 2001; Naghili, et al., 2013) and incubated overnight in ambient air at 35°C.

Colonies within each drop were counted. For drops containing 3 to 30 colonies, the cell density of the sample was calculated; if more than one dilution for a given sample had a countable number of colonies, the cell density of the two dilutions was averaged. If no drops had a countable number of colonies, the two drops above and below the countable range were averaged. The limit of detection was 300 CFU mL<sup>-1</sup>. Bactericidal activity was defined as a reduction of  $\geq 3 \log_{10}$  CFU mL<sup>-1</sup> at 24 hours compared to the starting inoculum (Leber, 2016; Pillai, et al., 2005).

## 2.5 Synergy studies

Checkboard synergy arrays consisting of orthogonal two-fold serial dilutions of antibiotic combinations were set up using the HPD300 as previously described (Brennan-Krohn, et al., 2017). Combinatorial fractional inhibitory concentrations  $\leq 0.5$  were considered synergistic;  $> 0.5$  and  $< 4$  were considered indifferent; and  $\geq 4$  were considered antagonistic (Odds, 2003).

## 2.6 Statistical Analysis

R was used to plot MIC distributions and for statistical analysis (RStudio, 2017; Team, 2017).  $P < 0.05$  was considered to be statistically significant.

## 3. Results

The *S. aureus* collection had a narrow apramycin MIC range of 4 to 32  $\mu\text{g/mL}$  (see Fig. 1). The MIC<sub>50</sub> and the MIC<sub>90</sub> were 8  $\mu\text{g/mL}$  and 16  $\mu\text{g/mL}$ , respectively. Based on the visual inspection method, an apramycin epidemiological cutoff value of 32  $\mu\text{g/mL}$  was assigned (Turnbridge & Patterson, 2007). A chi-square test of independence detected no significant

relationship between MSSA, MRSA and VISA strain phenotype and apramycin MIC distribution ( $P = 0.17$ ).

Apramycin exposure led to rapid rapid killing of clinical MSSA strain S27 and MRSA strain S19 at 1x-4x the broth microdilution MIC in time-kill experiments. However, regrowth was observed between 6 and 24 hours (Fig. 2). In contrast, VISA strain, FDA-CDC AR Bank #226, showed bacteriostatic activity at 1x-4x the broth microdilution MIC at all time points examined.

During checkerboard synergy testing of the same strains use for time-kill studies, both apramycin and gentamicin demonstrated indifference when tested in combination with vancomycin, daptomycin, and linezolid.

#### 4. Discussion

In our examination of human clinical isolates of *S. aureus*, we observed a very narrow distribution of MIC values. No isolate had an MIC above an epidemiological cutoff value of 32  $\mu\text{g/mL}$ , suggesting near to complete absence of apramycin modifying enzymes in our geographic region and in the smaller number of FDA-CDC VISA strains examined. This observation contrasted with a low prevalence of apramycin resistance in MRSA strains isolated from dairy cattle with mastitis and from diseased swine (Fessler, et al., 2011; Kadlec, et al., 2009). It is possible that selective pressure led to somewhat increased prevalence of apramycin resistance in these veterinary isolates. In the current study, we also found that apramycin demonstrated rapid, early, time-kill properties against MSSA and MRSA isolates with later regrowth, similar to previous reports for gentamicin (Schafer, et al., 2006), with static activity against a VISA isolate.

Apramycin is currently used as an orally administered, non-absorbable antibiotic to treat diarrheal diseases in poultry and livestock, as well as an intravenous treatment for pneumonia in calves and mastitis in cows and sheep (Livermore, et al., 2011; Ziv, et al., 1985; Ziv, et al., 1995). *S. aureus* is the most frequently isolated pathogen in bovine mastitis specimens (Bradley, et al., 2007), suggesting potential therapeutic efficacy against one type of *S. aureus* infection in



large mammals. In addition, efficacy of apramycin against a single MRSA strain was demonstrated in an immunocompromised murine septicemia model (Meyer, et al., 2014). Here, apramycin decreased bacterial burden in a dose-dependent manner by 2- to 3-log<sub>10</sub> in the blood and up to 4-log<sub>10</sub> in the kidneys. Therefore, there is evidence supporting therapeutic effect of apramycin against *Staphylococcal aureus* infection.

Notably, aminoglycosides are not used currently as single agents for treatment of *S. aureus* infections in humans based on limiting toxicities. Further, risk-benefit of short term use in combination with vancomycin and other agents for treatment of persistent bacteremia and prosthetic valve endocarditis, although recommended by some guidelines (Liu, et al., 2011) has been a matter of debate (Deresinski, 2009). It is possible that apramycin, based on a putatively more compelling side effect profile, may offer an alternative treatment for *S. aureus* infections as a single agent or in combination. However, basic pharmacokinetic and pharmacodynamic parameters still need to be defined in humans. Therefore use of apramycin for treatment of human *Staphylococcus aureus* infection as yet remains speculative.

## Conclusions

In this study, we found that apramycin shows consistent *in vitro* activity against contemporary *S. aureus* strains including MSSA, MRSA and VISA. Furthermore, apramycin demonstrated rapid bactericidal activity against MSSA and MRSA. Ultimate utility against human *S. aureus* infection will depend on pharmacokinetic and pharmacodynamic parameters that have yet to be fully established in animal models and investigated in humans. Nevertheless, the broad-spectrum *in vitro* activity of apramycin against multidrug-resistant *S. aureus* strains in combination with prior descriptions of activity against carbapenem-resistant Enterobacteriaceae, and multidrug-resistant *Acinetobacter baumannii* and *Pseudomonas aeruginosa*, suggests that further exploration of apramycin and/or derivatives as repurposed human therapeutics may be warranted.

## **Acknowledgements**

J.E.K. received support from a Chief Academic Officer's Pilot Award from Beth Israel Deaconess Medical Center. T.B.-K. and K.P.S. were supported by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under award numbers K08AI132716 and T32AI007061, and F32 AI124590, respectively. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

## **Disclosure Statement**

The HP D300 digital dispenser was provided by Tecan (Morrisville, NC). Tecan had no role in study design, data collection/interpretation, manuscript preparation, or decision to publish.

## References

- Akhi, MT, Ghotaslou, R, Alizadeh, N, Pirzadeh, T, Beheshtirouy, S, & Memar, MY (2017) High frequency of MRSA in surgical site infections and elevated vancomycin MIC. *Wound Medicine*, 17: 7-10. doi:10.1016/j.wndm.2017.01.002
- Bamberger, DM, & Boyd, SE (2005) Management of *Staphylococcus aureus* infections. *American Family Physician*, 72: 2474.
- Bradley, AJ, Leach, KA, Breen, JE, Green, LE, & Green, MJ (2007) Survey of the incidence and aetiology of mastitis on dairy farms in England and Wales. *Veterinary Record*, 160: 253-257.
- Brennan-Krohn, T, Truelson, KA, Smith, KP, & Kirby, JE (2017) Screening for synergistic activity of antimicrobial combinations against carbapenem-resistant Enterobacteriaceae using inkjet printer-based technology. *Journal of Antimicrobial Chemotherapy*, 72: 2775-2781. doi:10.1093/jac/dkx241
- Bruss, JB (2009) Lack of evidence associating nephrotoxicity with low-dose gentamicin for *Staphylococcus aureus* bacteremia and endocarditis. *Clinical Infectious Diseases*, 49: 806; author reply 807-808. doi:10.1086/605287
- Buchholtz, K, Larsen, CT, Hassager, C, & Bruun, NE (2009) Severity of gentamicin's nephrotoxic effect on patients with infective endocarditis: a prospective observational cohort study of 373 patients. *Clinical Infectious Diseases*, 48: 65-71. doi:10.1086/594122
- Buchholtz, K, Larsen, CT, Schaadt, B, Hassager, C, & Bruun, NE (2011) Once versus twice daily gentamicin dosing for infective endocarditis: a randomized clinical trial. *Cardiology*, 119: 65-71. doi:10.1159/000329842
- Chen, CY, Nace, GW, & Irwin, PL (2003) A 6 x 6 drop plate method for simultaneous colony counting and MPN enumeration of *Campylobacter jejuni*, *Listeria monocytogenes*, and *Escherichia coli*. *Journal of Microbiological Methods*, 55: 475-479.

Clinical and Laboratory Standards Institute (1999) *Methods for determining the bactericidal activity of antimicrobial agents; Approved guideline. CLSI document M-26A*. Wayne, PA: Clinical and Laboratory Standards Institute.

CLSI (2015) *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically; Approved Standard—10th Edition*. In *CLSI document M07-A10*. Wayne, PA: Clinical and Laboratory Standards Institute.

Cosgrove, SE, Vigliani, GA, Fowler, VG, Jr., Abrutyn, E, Corey, GR, Levine, DP, Rupp, ME, Chambers, HF, Karchmer, AW, & Boucher, HW (2009) Initial low-dose gentamicin for *Staphylococcus aureus* bacteremia and endocarditis is nephrotoxic. *Clinical Infectious Diseases*, 48: 713-721. doi:10.1086/597031

Deleo, FR, Otto, M, Kreiswirth, BN, & Chambers, HF (2010) Community-associated methicillin-resistant *Staphylococcus aureus*. *Lancet*, 375: 1557-1568. doi:10.1016/S0140-6736(09)61999-1

Deresinski, S (2009) Vancomycin in combination with other antibiotics for the treatment of serious methicillin-resistant *Staphylococcus aureus* infections. *Clinical Infectious Diseases*, 49: 1072-1079. doi:10.1086/605572

Dombrowski, JC, & Winston, LG (2008) Clinical failures of appropriately-treated methicillin-resistant *Staphylococcus aureus* infections. *Journal of Infection*, 57: 110-115. doi:10.1016/j.jinf.2008.04.003

Fessler, AT, Kadlec, K, & Schwarz, S (2011) Novel apramycin resistance gene *apma* in bovine and porcine methicillin-resistant *Staphylococcus aureus* ST398 isolates. *Antimicrobial Agents & Chemotherapy*, 55: 373.

Frippiat, F, Chandrikakumari, K, & Moutschen, M (2009) Gentamicin in infective endocarditis: how to use it? *Clinical Infectious Diseases*, 49: 320-321; author reply 321. doi:10.1086/600063

- Hawkins, C, Huang, J, Jin, N, Noskin, GA, Zembower, TR, & Bolon, M (2007) Persistent *Staphylococcus aureus* bacteremia: an analysis of risk factors and outcomes. *JAMA Intern Med*, 167: 1861.
- Hazlewood, KA, Brouse, SD, Pitcher, WD, & Hall, RG (2010) Vancomycin-associated nephrotoxicity: Grave concern or death by character assassination? *American Journal of Medicine*, 123: 182.e181-182.e187. doi:10.1016/j.amjmed.2009.05.031
- Herigstad, B, Hamilton, M, & Heersink, J (2001) How to optimize the drop plate method for enumerating bacteria. *J Microbiol Methods*, 44: 121-129.
- Jose, ML-N, Yaremi, Q, Laura, V, Ana, IM, & Francisco, JL-H (2010) New insights into the mechanism of aminoglycoside nephrotoxicity: an integrative point of view. *Kidney Int*, 79: 33. doi:10.1038/ki.2010.337
- Kadlec, K, Ehricht, R, Monecke, S, Steinacker, U, Kaspar, H, Mankertz, J, & Schwarz, S (2009) Diversity of antimicrobial resistance pheno- and genotypes of methicillin-resistant *Staphylococcus aureus* ST398 from diseased swine. *Journal of Antimicrobial Chemotherapy*, 64: 1156. doi:10.1093/jac/dkp350
- Kang, AD, Smith, KP, Berg, AH, Truelson, KA, Eliopoulos, GM, McCoy, C, & Kirby, JE (2018) Efficacy of apramycin against multidrug-resistant *Acinetobacter baumannii* in the murine neutropenic thigh model. *Antimicrobial Agents & Chemotherapy*, 62: e02585-02517.
- Kang, AD, Smith, KP, Eliopoulos, GM, Berg, AH, McCoy, C, & Kirby, JE (2017) *In vitro* apramycin activity against multidrug-resistant *Acinetobacter baumannii* and *Pseudomonas aeruginosa*. *Diagnostic Microbiology and Infectious Disease*, 88: 188-191. doi:10.1016/j.diagmicrobio.2017.03.006
- Leber, AL (2016) Time-Kill Assay for Determining Synergy. In A. L. Leber Ed, *Clinical Microbiology Procedures Handbook, Fourth Edition* (pp. 5.14.03.01-15.14.03.06). Washington, D.C.: ASM Press.

- Liu, C, Bayer, A, Cosgrove, SE, Daum, RS, Fridkin, SK, Gorwitz, RJ, Kaplan, SL, Karchmer, AW, Levine, DP, Murray, BE, M, JR, Talan, DA, Chambers, HF, & Infectious Diseases Society of, A (2011) Clinical practice guidelines by the infectious diseases society of america for the treatment of methicillin-resistant *Staphylococcus aureus* infections in adults and children. *Clinical Infectious Diseases*, 52: e18-55. doi:10.1093/cid/ciq146
- Livermore, DM, Mushtaq, S, Warner, M, Zhang, JC, Maharjan, S, Doumith, M, & Woodford, N (2011) Activity of aminoglycosides, including ACHN-490, against carbapenem-resistant Enterobacteriaceae isolates. *Journal of Antimicrobial Chemotherapy*, 66: 48-53. doi:10.1093/jac/dkq408
- Lodise, TP, Lomaestro, B, Graves, J, & Drusano, GL (2008) Larger vancomycin doses (at least four grams per day) are associated with an increased incidence of nephrotoxicity. *Antimicrobial Agents & Chemotherapy*, 52: 1330.
- Matt, T, Ng, CL, Lang, K, Sha, S-H, Akbergenov, R, Shcherbakov, D, Meyer, M, Duscha, S, Xie, J, Dubbaka, SR, Perez-Fernandez, D, Vasella, A, Ramakrishnan, V, Schacht, J, & Böttger, EC (2012) Dissociation of antibacterial activity and aminoglycoside ototoxicity in the 4-monosubstituted 2-deoxystreptamine apramycin. *Proceedings of the National Academy of Sciences, USA*, 109: 10984. doi:10.1073/pnas.1204073109
- Meyer, M, Freihofer, P, Scherman, M, Teague, J, Lenaerts, A, & Böttger, EC (2014) *In vivo* efficacy of apramycin in murine infection models. *Antimicrobial Agents & Chemotherapy*, 58: 6938. doi:10.1128/AAC.03239-14
- Naghili, H, Tajik, H, Mardani, K, Razavi Rouhani, SM, Ehsani, A, & Zare, P (2013) Validation of drop plate technique for bacterial enumeration by parametric and nonparametric tests. *Vet Res Forum*, 4: 179-183.
- Odds, FC (2003) Synergy, antagonism, and what the checkerboard puts between them. *Journal of Antimicrobial Chemotherapy*, 52: 1. doi:10.1093/jac/dkg301

- Pillai, SK, Moellering Jr, RC, & Eliopoulos, GM (2005) Antimicrobial Combinations. In V. Lorian Ed, *Antibiotics in Laboratory Medicine, Fifth Edition* (pp. 365-440). Philadelphia, PA: Williams and Wilkins.
- RStudio (2017) RStudio. In (1.0.143 ed.). Boston, MA.
- Schafer, JA, Hovde, LB, & Rotschafer, JC (2006) Consistent rates of kill of *Staphylococcus aureus* by gentamicin over a 6-fold clinical concentration range in an in vitro pharmacodynamic model (IVPDM). *Journal of Antimicrobial Chemotherapy*, 58: 108-111. doi:10.1093/jac/dkl216
- Smith, KP, & Kirby, JE (2016a) Evaluation of apramycin activity against carbapenem-resistant and -susceptible strains of Enterobacteriaceae. *Diagnostic Microbiology and Infectious Disease*, 86: 439-441. doi:10.1016/j.diagmicrobio.2016.09.002
- Smith, KP, & Kirby, JE (2016b) Validation of a high-throughput screening assay for identification of adjunctive and directly acting antimicrobials targeting carbapenem-resistant Enterobacteriaceae. *Assay and Drug Development Technologies*, 14: 194. doi:10.1089/adt.2016.701
- Smith, KP, & Kirby, JE (2016) Verification of an Automated, Digital Dispensing Platform for At-Will Broth Microdilution-Based Antimicrobial Susceptibility Testing. *Journal of Clinical Microbiology*, 54: 2288-2293. doi:10.1128/JCM.00932-16
- Steinkraus, G, White, R, & Friedrich, L (2007) Vancomycin MIC creep in non-vancomycin-intermediate *Staphylococcus aureus* (VISA), vancomycin-susceptible clinical methicillin-resistant *S. aureus* (MRSA) blood isolates from 2001–05. *Journal of Antimicrobial Chemotherapy*, 60: 788-794. doi:10.1093/jac/dkm258
- Tam, VH, Kabbara, S, Vo, G, Schilling, AN, & Coyle, EA (2006) Comparative pharmacodynamics of gentamicin against *Staphylococcus aureus* and *Pseudomonas aeruginosa*. *Antimicrobial Agents & Chemotherapy*, 50: 2626-2631. doi:10.1128/AAC.01165-05

Team, RC (2017) R: A Language and Environment for Statistical Computing. In. Vienna, Austria: R Foundation for Statistical Computing.

Tong, SYC, Davis, JS, Eichenberger, E, Holland, TL, & Fowler, VG (2015) *Staphylococcus aureus* infections: epidemiology, pathophysiology, clinical manifestations, and management. *Clinical Microbiology Reviews*, 28: 603. doi:10.1128/CMR.00134-14

Turnbridge, J, & Patterson, DL (2007) Setting and revising antibacterial susceptibility breakpoints. *Clinical Microbiology Reviews*, 20: 391. doi:10.1128/CMR.00047-06

Ziv, G, Bor, A, Sobagk, S, Elad, D, & Nouws, JFM (1985) Clinical pharmacology of apramycin in calves. *J Vet Pharmacol Ther*, 8: 95-104. doi:10.1111/j.1365-2885.1985.tb00930.x


Ziv, G, Kurtz, B, Risenberg, R, & Glickman, A (1995) Serum and milk concentrations of apramycin in lactating cows, ewes and goats. *J Vet Pharmacol Ther*, 18: 346-351. doi:10.1111/j.1365-2885.1995.tb00602.x




## Figures Legends

**Figure 1. Apramycin MIC distribution for *S. aureus* clinical isolates.** MSSA (methicillin-susceptible *S. aureus*); MRSA (methicillin-resistant *S. aureus*); VISA (vancomycin-intermediate *S. aureus*).


**Figure 2. Time-kill experiments.** Macrobroth time-kill analysis was performed against representative MSSA, MRSA, and VISA strains. Data points plotted at  $10^2$  cfu indicate no growth and correspond to the assay detection limit. Panel titles list the broth microdilution MIC associated with each isolate.



s27 MSSA (MIC = 16 µg/mL)



s19 MRSA (MIC = 8 µg/mL)



V7 VISA (MIC = 4 µg/mL)

